

symptoms than they would have had if the PDA had remained open.

Another important aspect is the accurate measurement of pulmonary vascular resistance.⁹ For patients breathing oxygen, it is crucial to include values for dissolved oxygen in the calculation of resistance. Failure to include dissolved oxygen in the measurement of pulmonary blood flow results in an underestimation of resistance. Many cardiac catheterisation laboratories do not have the facilities for the measurement of oxygen consumption,⁹ and instead, assumed values from tables are used. This inevitably provides an additional potential source of inaccuracy in pulmonary vascular resistance measurements. It is not clear from the text if assumed oxygen consumption was used or if resistance measurements took dissolved oxygen into account.

So, while on the one hand, it is possible to safely close a very large arterial duct in patients with pulmonary hypertension, it seems inevitable that at least some will develop progressive and fatal pulmonary vascular disease and eventually be worse-off than they would have been in terms of symptoms and life expectancy. Some patients had resistance values >15 units. The key to the safe management of these patients is the accurate measurement of pulmonary vascular resistance and management based on a resistance value rather than direct measurement of pulmonary artery pressure, which is never an accurate reflection of resistance. Management based on the direct measurement of pulmonary artery pressure during duct occlusion, although in common practice, is inevitably flawed.

Finally, in patients such as these, it is arguable that successful duct closure should be followed by an aggressive therapeutic approach to reduce pulmonary vascular resistance using pulmonary vasodilator agents, if possible.

Such treatment is of course extremely expensive, rarely successful in achieving a normal resistance and inappropriate in many developing countries, and should be part of research protocols and audit. For the group of patients reported, it is important that follow-up should include measurement of pulmonary vascular resistance at a given period after closure of the duct. Such a follow-up study might well be worthy of publication and will help to enhance our understanding of pulmonary vascular disease in this group of patients. However, the management of patients with a raised pulmonary vascular resistance and patent arterial duct must always be based on sound and reproducible physiological measurements and the underlying principle of "first do no harm."

Competing interests: None declared.

REFERENCES

- 1 **Gittenberger-De Groot AC.** Persistent ductus arteriosus: most probably a primary congenital malformation. *Br Heart J* 1977;**39**:610–8.
- 2 **Gross RE, Hubbard JP.** Surgical ligation of a patent ductus arteriosus. A report of first successful case. *JAMA* 1930;**112**:729–31.
- 3 **Portsmann W, Wierny L, Warnke H, et al.** Catheter closure of patent ductus arteriosus, 62 cases treated without thoracotomy. *Radiol Clin North Am* 1971;**9**:203–18.
- 4 **Rashkind WJ, Mullins CE, Hellenbrand WE, et al.** Non-surgical closure of patent ductus arteriosus: clinical application of the Rashkind PDA Occluder System. *Circulation* 1987;**75**:583–92.
- 5 **Hosking MC, Benson LN, Musewe N, et al.** Transcatheter occlusion of the persistently patent ductus arteriosus. *Circulation* 1991;**84**:2313–17.

- 6 **Jones JC.** Twenty-five years experience with the surgery of patent ductus arteriosus. *J Thorac Cardiovasc Surg* 1965;**50**:149–65.
- 7 **Campbell M.** Natural history of persistent ductus arteriosus. *Br Heart J* 1968;**30**:4.
- 8 **Yan C, Zhao S, Jiang S, et al.** Transcatheter closure of patent ductus arteriosus with severe pulmonary arterial hypertension in adults. *Heart* 2007;**93**:514–18.
- 9 **Bush A, Busst CM, Haworth SG, et al.** Correlations of lung morphology, pulmonary vascular resistance, and outcome in children with congenital heart disease. *Br Heart J* 1988;**59**:480–6.
- 10 **Heath D, Edwards JE.** The pathology of hypertensive pulmonary vascular disease: a description of six grades of structural changes in the pulmonary arteries with special reference to congenital cardiac septal defects. *Circulation* 1958;**18**:533–43.

WEB TOP 10

www.heartjnl.com

These articles scored the most hits on *Heart's* website during January 2007

1 The value of risk scores

JS Jürgensen

December 2006;**92**:1713–4. (Editorial)

2 JBS 2: Joint British Societies' guidelines on prevention of cardiovascular disease in clinical practice

December 2005;**91**(suppl V):1–52. (Supplement)

3 Antiplatelet treatment for coronary heart disease

N Clappers, MA Brouwer, FWA Verheugt

February 2007;**93**:258–65. (Education in Heart)

4 Antimicrobial prophylaxis for endocarditis: emotion or science?

H Ashrafian, RG Bogle

January 2007;**93**:5–6. (Editorial)

5 Diagnosis and treatment of syncope

M Brignole

January 2007;**93**:130–6. (Education in Heart)

6 Heart failure with a normal ejection fraction

JE Sanderson

February 2007;**93**:155–8. (Heart review)

7 Cardiovascular, cerebrovascular, and respiratory changes induced by different types of music in musicians and non-musicians: the importance of silence

L Bernardi, C Porta, P Sleight

April 2006;**92**:445–2. (Cardiovascular medicine)

8 Management of tricuspid valve regurgitation

MJ Antunes, JB Barlow

February 2007;**93**:271–6. (Education in Heart)

9 Drug induced QT prolongation and torsades de pointes

YG Yap, AJ Camm

November 2003;**89**:1363–72. (Education in Heart)

10 Thrombosis of prosthetic heart valves: diagnosis and therapeutic considerations

R Roudaut, K Serri, S Lafitte

January 2007;**93**:137–42. (Education in Heart)

Visit the *Heart* website for hyperlinks to these articles, by clicking on "Top 10 papers"

www.heartjnl.com